

Kilohertz QPOs in Neutron Star Binaries modeled as Keplerian Oscillations in a Rotating Frame of Reference

Vladimir Osherovich

NASA/Goddard Space Flight Center/RITSS, Greenbelt MD 20771 USA;

vladimir@urap.gsfc.nasa.gov

Lev Titarchuk

NASA/Goddard Space Flight Center, Greenbelt MD 20771, and George Mason

University/CSI, USA; titarchuk@lheavx.gsfc.nasa.gov

Received _____; accepted _____

ABSTRACT

Since the discovery of kHz quasi-periodic oscillations (QPO) in neutron star binaries, the difference between peak frequencies of two modes in the upper part of the spectrum, i.e. $\Delta\omega = \omega_h - \omega_k$ has been studied extensively. The idea that $\Delta\omega$ is constant and (as a beat frequency) is related to the rotational frequency of the neutron star has been tested previously. The observed decrease of $\Delta\omega$ when ω_h and ω_k increase has weakened the beat frequency interpretation. We put forward a different paradigm: a Keplerian oscillator under the influence of the Coriolis force. For such an oscillator, ω_h and the assumed Keplerian frequency ω_k hold an upper hybrid frequency relation: $\omega_h^2 - \omega_k^2 = 4\Omega^2$, where Ω is the rotational frequency of the star’s magnetosphere near the equatorial plane. For three sources (Sco X-1, 4U 1608-52 and 4U 1702-429), we demonstrate that the solid body rotation $\Omega = \Omega_0 = \text{const.}$ is a good first order approximation. Within the second order approximation, the slow variation of Ω as a function of ω_K reveals the structure of the magnetospheric differential rotation. For Sco X-1, the QPO have frequencies ~ 45 and 90 Hz which we interpret as the 1st and 2nd harmonics of the lower branch of the Keplerian oscillations for the rotator with $\mathbf{\Omega}$ not aligned with the normal of the disk: $\omega_L/2\pi = (\Omega/\pi)(\omega_K/\omega_h) \sin \delta$ where δ is the angle between $\mathbf{\Omega}$ and the vector normal to the disk.

Subject headings: accretion, accretion disks — radiation mechanisms: thermal —stars:neutron — X-rays: general

1. Introduction

Kilohertz quasi-periodic oscillations (KQPO) have been discovered by the Rossi X-ray Timing Explorer (RXTE) in a number of low mass X-ray binaries (Strohmayer *et al.* 1996). The existence of two observed peaks with frequencies ω_K and ω_h in the upper part of the QPO spectrum became a natural starting point in modeling the phenomenon. Attempts have been made to relate ω_K , ω_h and the peak difference frequency $\Delta\omega = \omega_h - \omega_K$ with the neutron star spin and possible Keplerian motion of the hot matter surrounding the star [see origin of Keplerian motion discussed in Titarchuk, Lapidus & Muslimov 1998 (hereafter TLM) and Titarchuk & Osherovich 1999].

In the beat frequency model, the burst oscillations and the kHz peak separation $\Delta\omega$ are both considered to be close to the neutron star spin frequency and thus $\Delta\omega$ is predicted to be constant (see, e.g., review by van der Klis 1998). However, recent observations of kHz QPO's in Sco X-1, 4U 1608-52 and 4U 1702-429 showed that $\Delta\omega$ decreases systematically when ω_K and ω_h increases (van der Klis et al. 1997; Mendez et al. 1998; Markwardt, Strohmayer & Swank 1999). Indications of such a decrease was found by Wijnands et al. (1998) and Ford et al. (1998) for 4U 1735-44 as well. Psaltis et al. (1998) showed that the measured $\Delta\omega$ in all other known sources is consistent with the values found in Sco X-1 and 4U 1608-52. These sources have become standards for comparison between the kHz QPO phenomena in different sources because of the high accuracy of measurements and the wide range of values of ω_K and ω_h available. For Sco X-1 and 4U 1608-52, $\Delta\omega/2\pi$ changes from 320 Hz to 220 Hz when $\omega_K/2\pi$ changes from 500 Hz to 850 Hz

In the lower part of the Sco X-1 spectra, van der Klis et al. (1997) found two frequencies ~ 45 Hz (referred to below as $\omega_L/2\pi$) and 90 Hz which also slowly increase with increase of ω_K and ω_h ; presumably 90 Hz is the second harmonic of $\omega_L/2\pi$. Any consistent model faces a challenging task of describing the dependence of $\Delta\omega$ and ω_L on ω_K and ω_h for the

available sources.

In the beat-frequency model (BFM) considered by Miller, Lamb & Psaltis (1998), the kHz QPO's are identified with the orbital frequency at the sonic radius and its beat with the neutron star spin, and the lower frequency of horizontal branch oscillations (HBO) in Z-sources with the beat of the orbital motion at the magnetospheric radius with the neutron star spin. In the general-relativistic (GR) precession/apsidal motion discussed by Stella and Vietri (1998,1999), the kHz peaks are due to orbital motion and GR apsidal motion of a slightly eccentric orbit, and low-frequency QPO to the Lense-Thirring precession of the same orbit. Titarchuk and Muslimov (1997) suggested that a set of peaks seen in the kHz QPOs can be naturally explained in terms of the effect of rotational splitting of the main oscillation frequency in the disk. The type of peak (eigenmodes) is dependent on mode numbers and a parameter describing disk structure. In TLM the authors argue that this effect of the rotational splitting originated in a disk region identified as the centrifugal barrier (CB) region oscillated in the vertical and radial directions. The size oscillations of the CB disk region and reprocessing of the disk photons there (due to Comptonization) led to an oscillation signal.

In this *Letter*, we take a different approach and we suggest that a kHz QPO can be explained as oscillations of large scale inhomogeneities (hot blobs) thrown into the neutron star's magnetosphere. Participating in the radial oscillations with Keplerian frequency ω_K , such blobs are simultaneously under the influence of the Coriolis force. The present model is very different than the TLM one where the site of the rotational splitting was associated with the CB disk region. Thus, studying the Keplerian oscillator in the rotating frame of reference (magnetosphere), we attempt to relate $\Delta\omega$ and ω_L with ω_K and ω_h . In our model, we compare results of linear theory with the observations for three sources. We stress the fundamental role of the Keplerian frequency (which we identify with ω_K) and the upper

hybrid frequency relation between ω_K and ω_h . In this first *Letter*, we depict the main features of the kHz QPO phenomena. The discussion identifies some limitations of our approach and suggests further directions of the research we shall pursue.

2. Keplerian Oscillator Under the Influence of the Coriolis Force

We assume that the interplay between the centrifugal, gravitational force and magnetic force maintains radial oscillations (along the axis x) with the Keplerian frequency

$$\omega_K = \left(\frac{GM}{R^3} \right)^{1/2} \quad (1)$$

where G is the gravitational constant, M is the mass of the compact object and R is the radius of an orbit. Self-similar spheromak-type models of a magnetic atmosphere can serve as an example of the exact time dependent MHD solutions in which non-radial equilibrium is maintained by magnetic force, while all motion is strictly radial (Farrugia et al. 1995 and references there in). Taking the magnetospheric rotation with angular velocity Ω (not perpendicular to the plane of the equatorial disk), the small amplitude oscillations of such a blob thrown into the magnetosphere are described by equations (Landau and Lifshitz, 1960)

$$\ddot{x} + \omega_K^2 x - 2\dot{y}\Omega \cos \delta = 0 \quad (2)$$

$$\ddot{y} + 2\dot{x}\Omega \cos \delta - 2\dot{z}\Omega \sin \delta = 0 \quad (3)$$

$$\ddot{z} + 2\dot{y}\Omega \sin \delta = 0 \quad (4)$$

where δ is the angle between Ω and the vector normal to the plane of radial oscillations, y is the azimuthal component of the displacement vector and z is the vertical component which is perpendicular to the plane of Keplerian oscillations. The dispersion relation for the frequency ω

$$\omega^2[\omega^4 - (\omega_K^2 + 4\Omega^2)\omega^2 + 4\Omega^2\omega_K^2 \sin^2 \delta] = 0 \quad (5)$$

besides the non-oscillating mode ($\omega = 0$), describes two spectral branches (eigenmodes), high and low:

$$(\omega^2)_{1,2} = \frac{\omega_h^2 \pm (\omega_h^4 - 16\Omega^2\omega_K^2 \sin^2 \delta)^{1/2}}{2} \quad (6)$$

where

$$\omega_h = (\omega_K^2 + 4\Omega^2)^{1/2} \quad (7)$$

is the analog of the upper hybrid frequency f_{uh} in plasma physics (Akhiezer et al., 1975; Benson 1977). When the angle δ is small (as we believe is indeed our case):

$$\omega_1 = \omega_h \quad (8)$$

is related with the radial eigenmode and

$$\omega_2 \equiv \omega_L = 2\Omega(\omega_K/\omega_h) \sin \delta \quad (9)$$

is related with the vertical eigenmode. For $\delta \ll 1$, corrections for ω_1 are of the second order in δ , but for ω_2 , the angle dependent term is the main term.

3. Comparison of the Model with Observations

According to formula (7)

$$\Omega = \frac{(\omega_h^2 - \omega_K^2)^{1/2}}{2} \quad (10)$$

is the rotational frequency, which, for the highly conductive plasma in the case of corotation, should be approximately constant ($\Omega = \Omega_0$). Interpreting the two frequencies in the upper part of the kHz spectrum as ω_K and ω_h [where $\omega_K < \omega_h$] for Sco X-1, we plotted in Figure 1, $\nu = \Omega/2\pi$ versus $\nu_K = \omega_K/2\pi$ (solid circles). Indeed, one may notice that as a first approximation $\Omega = \Omega_0 = \text{const}$. The slow (but systematic) variation of $\Omega/2\pi$ is between 330 and 350 Hz which should be compared with the variation by a factor of 1.5 for $\Delta\omega = \omega_h - \omega_K$ for the same range of ω_K . If the magnetosphere corotates with the neutron

star (solid body rotation), then our procedure would determine the spin rotation of the star. In fact, there is a differential rotation profile which depends on the magnetic field $\mathbf{B}$ which can be presented in terms of the Chandrasekhar potential A (Chandrasekhar, 1956)

$$\mathbf{B} = \frac{1}{R} \left(-\frac{1}{R} \frac{\partial A}{\partial \mu}, -\frac{1}{(1-\mu^2)^{1/2}} \frac{\partial A}{\partial R}, \frac{B^*(A)}{(1-\mu^2)^{1/2}} \right) \quad (11)$$

where $\mu = \cos \theta$, θ is the colatitude in the spherical system of coordinates and $B^*(A)$ is a function of A only. For a combination of dipole, quadrupole and octupole fields

$$A = A_0(1 - \mu^2) \left[\frac{1}{R} + \frac{q}{R^2} \mu - \frac{b}{R^3} (5\mu^2 + 2) \right] \quad (12)$$

where A_0 , q and b are constants. For a highly conductive plasma, the MHD induction equation leads to the iso-rotation theorem of Ferraro (see Alfvén and Falthammar, 1963) which in our notation means that Ω is a function of A only. Following a recent suggestion of Osherovich and Gliner (1999), we assume that in the vicinity of a magnetic star $\Omega(A)$ has a maximum and therefore can be presented as

$$\Omega(A) = \Omega_0 - a^2 A^2 \quad (13)$$

where Ω_0 and a are constants. Since in the equatorial plane ($\mu = 0$), according to (12), the quadrupole term does not contribute to A , from formula (13) we find

$$\Omega(A) = \Omega_0 - \left(\frac{\alpha'}{R} - \frac{\beta'}{R^3} \right)^2 \quad (14)$$

where α' and β' are positive constants. Expressing R through the Keplerian frequency $\nu_K = \omega_K/(2\pi)$ from equation (1), we find that formula (14) leads to the following presentation for Ω :

$$\Omega(\nu_K)/2\pi = C_0 + C_1 \nu_K^{4/3} + C_2 \nu_K^{8/3} + C_3 \nu_K^4 \quad (15)$$

where $C_2 = 2\sqrt{C_1 C_3}$ (not a free parameter because the radial part of the expansion is a full square of $\nu_K^{2/3}$ and ν_K^2 with new positive constants α and β). The fit for Sco X-1 (solid line

in Figure 1) by formula (16) demonstrates the agreement of the proposed model with the data. Indeed, for Sco X-1, we find $C_0 = \Omega_0/2\pi = 345$ Hz, $C_1 = -\alpha^2 = -3.29 \cdot 10^{-2}$ Hz $^{-1/3}$, $C_2 = 2\alpha\beta = 1.017 \cdot 10^{-5}$ Hz $^{-5/3}$ and $C_3 = -\beta^2 = -7.76 \cdot 10^{-10}$ Hz $^{-3}$ with high $\chi^2 = 37.6/39$.

In our fitting procedure, C_0 , C_1 , C_2 and C_3 are treated as four free parameters, i.e., the relation between C_2 and the two constants C_1 and C_3 is not enforced. The fact that the derived C_2 indeed satisfies the relation $C_2 = 2\sqrt{C_1 C_3}$ shows the internal consistency of the model.

Three available data points for the 4U 1702-429 source are consistent with a constant value of rotation frequency $\Omega_0/2\pi = 380$ Hz, but the small number of data points does not allow us to recover the profile of $\Omega(\nu_K)$. However, the fit of 4U 1608-52, shown in Figure 2, demonstrates that the excellent agreement between our model and observations for Sco X-1 is not accidental. For 4U 1608-52, we found $C_0 = \Omega_0/2\pi = 345$ Hz, $C_1 = -3.91 \cdot 10^{-2}$ Hz $^{1/3}$, $C_2 = 1.23 \cdot 10^{-5}$ Hz $^{-5/3}$ and $C_3 = -9.45 \cdot 10^{-10}$ Hz $^{-3}$ with $\chi^2 = 12.02/13$.

Sco X-1 allows us to check the prediction of the model for ω_L [formula (10)]. Taking the profile $\Omega(\nu_K)$, presented in Figure 1, we plot $\nu_L = \omega_L/2\pi$ in Figure 3. Choosing $\delta = 5.5^\circ$, we found that our plot for ν_L and $2\nu_L$ fit the data for the observed frequencies of 45 Hz and 90 Hz, respectively. From these results, we conclude that the rotational axis of Sco X-1 and the normal to the plane of radial oscillations do not coincide but are separated by 5.5° .

4. Summary and Discussion

The main thrust of this work is based on the hypothesis that two of the observed frequencies in the upper part of the kHz QPO spectrum are the Keplerian frequency (ω_K) and the upper hybrid frequency (ω_h) of the Keplerian oscillator in the frame of reference rotating with the angular frequency Ω . For three sources, we showed that

Ω calculated according to the formula $\Omega = \sqrt{\omega_h^2 - \omega_K^2}/2$ changes with ω_K significantly less than $\Delta\omega = \omega_h - \omega_K$, suggesting approximately constant rotation of the star's magnetosphere. The detailed profile of Ω is successfully modeled as a function of ν_K within the dipole-quadrupole-octupole approximation of the magnetic field for Sco X-1 and for 4U 1608-52. For Sco X-1, the derived $\Omega(\nu_K)$ is used to compare predictions of the model for the lower branch frequency $\omega_L/2\pi$ with observations of 45 Hz and 90 Hz (presumably $2\omega_L/2\pi$). The last fit is done with only one unknown parameter δ (angle between the $\mathbf{\Omega}$ and the normal vector to the plane of Keplerian oscillations).

A good fit for ω_L and $2\omega_L$ confirmed the suggested model and revealed the expectedly small angle $\delta = 5.5^\circ$. The profile for $\Omega(\nu_K)$ has been modeled strictly in the equatorial plane ($\mu = 0$) for the ideally aligned rotator ($\delta = 0$). With $\delta \neq 0$, the next approximation will reveal the asymmetry of the QPO spectrum due to the quadrupole term. It has been shown (Osherovich, Tzur and Gliner, 1984) for the solar corona that the quadrupole term introduces North-South asymmetry in the makeup of the magnetic atmosphere surrounding a gravitating object. We shall pursue a similar study for the neutron star with the expectation of finding a signature of such asymmetry in the kHz QPO. Recent measurements show that the octupole field is comparable to the global dipole field near the sun, while the quadrupole field contributes 20 – 30% of the total field (Wang et al., 1997; Osherovich et al., 1999).

In this *Letter*, we restrict the scope of the work to a model for the linear Keplerian oscillator. The presence of the second harmonic (90 Hz) of the 45 Hz mode, noticed by van der Klis et al. (1997), strongly suggests that a weakly nonlinear model is desirable. Some oscillations below ω_L (for Sco X-1 oscillations with frequencies 10 – 20 Hz) we attribute to the innermost edge of the Keplerian disk, which adjusts itself to the rotating central object (i.e., neutron star). The physics of these oscillations related to accreting matter with

a large angular momentum is outside the scope of this Letter and it have been described in a separate paper (Titarchuk & Osherovich 1999).

We are grateful to J. Fainberg and R.F. Benson for help in the preparation of this paper and discussions and for fruitful suggestions by the referee. L.T. thanks NASA for support under grants NAS-5-32484 and RXTE Guest Observing Program and Jean Swank for support of this work. In particular, we are grateful to Michiel van der Klis and Mariano Mendez for the data which enabled us to make detailed comparisons of results of our model with the data.

REFERENCES

- Akhiezer, A.I., Akhiezer, A.I., Polovin, R.V., Sitenko, A.G. & Stepanov, K.N. 1975, Plasma Electrodynamics (New York: Pergamon)
- Alfvén, H., & Falthammar, C-G. 1963, Cosmical Electrodynamics. Fundamental Principles (London: Qxford University Press)
- Benson, R.F. 1977, Rad. Sci., 12, 861
- Chandrasekhar, S. 1956, Proc. Acad. Sci. 42, 1
- Farrugia, C.J., Osherovich, V.A. & Burlaga, L.F. 1995, J. Geophys. Res., 100, 293
- Ford, E., van der Klis, M., van Paradijs, J., Wijnands, R., Mendez, M., & Kaaret, P. 1998, ApJ, 508, L155
- Landau, L., & Lifshitz, E. 1960, Mechanics (New York: Pergamon)
- Markwardt, C.B., Strohmayer, T.E., & Swank, J.H. 1999, ApJ, in press
- Mendez, M., van der Klis, M., Ford, E.C., Wijnands, R., van Paradijs, J., & Vaughan, B.A. 1998, ApJ, 505, L23
- Miller, M.C., Lamb, F.K. & Psaltis, D. 1998, ApJ, 508, 791
- Osherovich, V.A., Tzur, I., & Gliner, E.B. 1984, ApJ, 284, 412
- Osherovich, V.A., & Gliner, E.B. 1999, in preparation
- Psaltis, D., et al. 1998, ApJ, 501, L95
- Swank J.H., 1998, IAU SYMP. 188, 107
- Shakura, N. I. & Sunyaev, R. A. 1973, A&A, 24, 337

- Stella, L., & Vietri, M, 1998, ApJ, 492, L59
- Stella, L., & Vietri, M. 1999, Phys. Rev. Letters, 82, 17
- Strohmayer, T. E., Zhang, W., Swank, J. H., Smale, A., Titarchuk, L., Day, C., & Lee, U. 1996, ApJ, 469, L9
- Swank, J., astro-ph/9802188 to appear in “The active X-ray Sky” eds. L. Scarsi, H Bradt, P. Giommi, and F. Fiore (1998)
- Titarchuk, L., Lapidus, I.I.,& Muslimov, A. 1998, ApJ, 499, 315
- Titarchuk, L.,& Osherovich, V. 1999, ApJ, 518, in press
- Titarchuk, L., & Muslimov, A. 1997, A&A, 323, L5
- van der Klis, M. 1998, in AIP Conf Proc 431, 361
- van der Klis, M., Wijnands, R.A.D., Horne, K. & Chen, W. 1997, ApJ, 481, L97
- Wijnands, R.A.D., & van der Klis, M. 1997, Nature, 394, 344
- Wang, Y-M., *et al.* 1997, ApJ, 485, 875

Fig. 1.— The inferred rotational frequency of the NS magnetosphere (see Eq. 10) as a function of the frequency of the low kHz QPO peak (Keplerian frequency). For Sco X-1 (van der Klis et al. 1997) (solid circles) and for 4U 1702-429 (Markwardt et al. 1999) (open squares). The solid (in the case of Sco X-1) line and dashed line (in the case of 4U 1702-429) are theoretical curves calculated using the multipole expansion of the rotational frequency (Eq.15).

Fig. 2.— The inferred rotational frequency of the NS magnetosphere (see Eq. 10) as a function of the frequency of the low kHz QPO peak (Keplerian frequency). For 4U 1608 (Mendez et al. 1998) (solid squares). The solid line is the theoretical curve (see Eq.15).

Fig. 3.— HBO frequency vs high peak KQPO frequency. for Sco X-1 (van der Klis et al. 1997) (solid circles). The solid line is theoretical dependence of the low branch oscillation frequencies (40-50 Hz) and second harmonics (80-100 Hz) on the high hybrid frequencies, calculated using the multipole expansion of the rotational frequency (Eq. 16) with $\delta = 5.5^\circ$.





